

Budget Impact Analysis (BIA)

Esempio di BIA correlata all'aderenza terapeutica: esperienza sui DOAC nella FANV

Dati di aderenza

	Total	Apixaban	Dabigatran	Rivaroxaban
Adh mean ± sd	0,85±0,15	0,86±0,13	0,72±0,22	0,88±0,11
Adh ≥ 0,8 N (%)	3480(81)	1359(84)	276(49)	1847(89)
Adh < 0,8 N (%)	790(19)	264(16)	287(51)	237(11)
Adh 1Y mean ± sd	0,83±0,18	0,84±0,17	0,71±0,24	0,86±0,15
Adh 2Y mean ± sd	0,85±0,15	0,86±0,14	0,72±0,22	0,87±0,13
Adh 3Y mean ± sd	0,85±0,15	0,86±0,13	0,72±0,22	0,87±0,12



Adherence, persistence and switching rates of apixaban, dabigatran and rivaroxaban in non-valvular atrial fibrillation: a multicentre real-life analysis at 3 years

Alessia Romagnoli ,¹ Fiorenzo Santoleri,² Alberto Costantini,² Angelora Di Risio¹

Input da considerare

- Costo dei farmaci
- % di ricoveri correlati all'aderenza:


Increased Risk of Stroke Due to Non-adherence and Non-persistence with Direct Oral Anticoagulants (DOACs): Real-World Analyses Using a Nested Case–Control Study from The Netherlands, Italy and Germany

Emily Holthuis¹ · Elisabeth Smits¹ · George Spentzouris² · Dominik Beier³ · Dirk Enders³ · Rosa Gini⁴ · Claudia Bartolini⁴ · Giampiero Mazzaglia⁵ · Fernie Penning-van Beest¹ · Ron Herings^{1,6}

- Aderenza ≥ 0,8: 44% ricoveri per stroke
- Aderenza < 0,8: 56% ricoveri per stroke



Costo medio annuale ricoveri per stroke:



Totale: € 1.567 Ospedalizzazione



Risultati

